

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## DIOPHANTINE ANALYSIS.

## 101. Proposed by HARRY S. VANDIVER, Bala, Pa.

Prove that it is impossible to find integral values for x, y, and z such that the relation  $x^2y + xz^2 = y^2z$  is satisfied.

## II. Solution by W. F. KING, Ottawa, Canada.

We may assume that x, y, and z have no factor common to them all. For if they have a common factor n, each term of the equation  $x^2y+xz^2=y^2z$  may be divided by  $n^3$ , and there will then be left an equation of the same form, in which x, y, and z have no common factor.

Let the greatest common measure of x and y be a; of y and z, b; and of z and x, c. Then we shall have

$$x=cax_1,$$
  
 $y=aby_1,$   
 $z=bcz_1.$ 

Now observe that since b is the greatest common measure of y and z,  $ay_1$  is prime to  $cz_1$ ; similarly is  $bz_1$  to  $ax_1$ , and  $cx_1$  to  $by_1$ .

Hence a, b, and c are all prime to one another; so are  $x_1$ ,  $y_1$ , and  $z_1$  to one another. Also a is prime to  $z_1$ , b to  $x_1$ , and c to  $y_1$ . Substituting in the equation the above values of x, y, and z, and dividing by abc, we have

$$a^2 c x_1^2 y_1 + c^2 b z_1^2 x_1 = b^2 a y_1^2 z_1$$
.

Divide by 
$$x_1$$
; then  $a^2c x_1y_1 + c^2b z_1^2 = \frac{b^2a y_1^2z_1}{x_1}$ .

The left hand side is integral, therefore  $b^2a y_1^2z_1$  is divisible by  $x_1$  But as above shown,  $x_1$  is prime to b,  $y_1$  and  $z_1$ . Hence  $a/x_1$  must be an integer.

Again, divide the above equation by a. Then

$$acx_1^2y_1 + \frac{c^2bz_1^2x_1}{a} = b^2y_1^2z_1$$
.

Hence, as before,  $\frac{c^2b \ z_1^2 x_1}{a}$  is an integer. But a is prime to b, c, and  $z_1$ . Therefore,  $x_1/a$  is an integer. Now since  $a/x_1$  and  $x_1/a$  are both integers,  $x_1=a$ . Similarly,  $y_1=b$ ,  $z_1=c$ . Therefore the equation

$$a^2c x_1^2y_1 + c^2b z_1^2x_1 = b^2a y_1^2z_1$$

becomes  $a^4bc+abc^4=ab^4c$ , or dividing by abc,  $a^3+c^3=b^3$ , a known impossible form.